



Knowledge grows

YaraVita™ BRASSITREL PRO

A formulated multi-nutrient foliar treatment for brassicas, legumes and other vegetable and arable crops

Guaranteed Analysis

Nitrogen (N)	6.9%	Boron (B)	6.0%
Calcium (Ca)	8.9%	Manganese (Mn)	7.1%
Magnesium (Mg)	7.1%	Molybdenum (Mo)	0.5%

Why Foliar Apply?

Foliar sprays ensure precise application of the right nutrient mix at the right time, and can be specifically targeted to the leaf or fruit, to suit an immediate crop need.

Foliar application also provides nutrients for immediate uptake by the leaves or fruits. As a result, the grower is not reliant on the right soil, pH or growing media conditions and can quickly put the crop back on course.

Magnesium, as well as the micronutrients boron, manganese and molybdenum are essential for crop establishment as well as winter hardiness of autumn sown crops. Stem extension is a period of high nutrient demand and often uptake of nutrients from the soil cannot keep pace with demand.

BRASSITREL PRO is formulated to supply all these vital nutrients in a concentrated, safe and balanced way at key moments in the growing season.



Benefits:

- Formulated for safe application at critical growth stages to satisfy crop requirements
- Widely tank mixable with other crop sprays. Visit www.tankmix.com for details
- Proven, reliable performance
Trialed and tested on a wide range of crops around the world
- High quality, consistent products. Manufactured to ISO 9001 quality assurance standards
- Easy to use flowable formulation.
- Pours and disperses easily and quickly into the spray tank
- High nutrient content means lower application rates reducing handling time and waste packaging.
- Manufactured to ISO 9001 quality assurance standards.



Application rates and timings

Beans, Legumes/Pulses, Peas, Soya Bean: 3 to 4 l/ha as early in the growing season as possible, provided there is adequate leaf cover to intercept the spray (e.g. 4 to 6 leaf stage). Repeat as necessary at 10 to 14 day intervals up to one month before harvest. N.B. Do not spray during flowering. Water rate: 200 l/ha.

Brassicas, Broccoli, Brussel Sprouts, Cauliflower, Kale, Lettuce (Field Grown), Radish (Field Grown) Spinach, Swede, Turnip: 3 to 4 l/ha as early in the growing season as possible, provided there is adequate leaf cover to intercept the spray (e.g. 4 to 6 leaf stage). Repeat as necessary at 10 to 14 day intervals up to one month before harvest. Water rate: 200 l/ha.

Carrot: Two applications of 3 l/ha applied when the crop is 15 cm tall and again 2 weeks later. Water rate: 200 l/ha.

Cereals: 3 l/ha applied at any time from the 2 leaf stage to 1st node detectable (Zadoks G.S. 12 to 31). Water rate: 200 l/ha.

Oilseed Rape: 3 to 4 l/ha at the 4 to 9 leaf stage and again at the onset of the rapid stem extension phase up to the start of flowering. Repeat at 10 to 14 day intervals within these growth stages as necessary. Water rate: 50 to 200 l/ha.

Onion: 2 applications of 3 l/ha applied as soon as there is adequate leaf cover and repeated 2 weeks later. Water rate: 200 l/ha.

Potatoes: 3 l/ha applied one week after 100 % emergence with a second application 10 to 14 days later. Water rate: 200 l/ha.

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS
Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes (temperatures exceeding 40°C). Dispose of contents and container in accordance with all local, regional, national and international regulations

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.

Yara Fertilizers (NZ) Ltd

Tel: +64 6 8776600

Email: nz.enquiries@yara.com

Website: www.yara.co.nz

