



Knowledge grows

YaraVita™ CROPLIFT K

A formulated multi-nutrient product for foliar application

Guaranteed Analysis

Nitrogen (N)	5%	Manganese (Mn)	0.26%
Phosphorus (P)	4.4%	Copper (Cu)	0.2%
Potassium (K)	29%	Zinc (Zn)	0.14%
Sulphur (S)	4%	Boron (B)	0.04%
Magnesium (Mg)	1.2%	Iron (Fe)	0.02%



Why Foliar Apply?

Foliar sprays ensure precise application of the right nutrient mix at the right time, and can be specifically targeted to the leaf or fruit, to suit an immediate crop need.

Foliar application also provides nutrients for immediate uptake by the leaves or fruits. As a result, the grower is not reliant on the right soil, pH or growing media conditions and can quickly put the crop back on course

YaraVita CROPLIFT K is a foliar fertiliser containing multiple nutrients and micronutrients for foliar application to a wide range of crops during times of stress or periods of rapid growth.

Benefits:

- Formulated for safe application at critical growth stages to satisfy crop requirements.
- Widely tank mixable with other crop sprays. Visit www.tankmix.com for details.
- Proven, reliable performance. Trialled and tested on a wide range of crops around the world
- High nutrient content means lower application rates reducing handling time and waste packaging.
- Easy to use soluble powder. Pours and disperses easily and quickly into the spray tank
- High quality, consistent products.
- Manufactured to ISO 9001 quality assurance standards.

Yara Fertilizers (NZ) Ltd

Tel: +64 6 8776600

Email: nz.enquiries@yara.com

Website: www.yara.co.nz



Application rates and timings

Foliar Application

All crops (field grown) except those specified below: 2.5 to 5 kg/ha. Repeat at 10 to 14 day intervals as necessary. Avoid applications during flowering. Also, apply as necessary to perennial fruit crops post-harvest. Water rate: 200 l/ha (500 to 1000 l/ha for fruit crops). Alternatively: Spray to run off using a concentration of 2.5 grammes per litre. Allow 10 to 14 days between successive applications.

Apple: 5 kg/ha at 10 to 14 day intervals as required commencing 4 weeks after petal fall. Water rate: 500-1000 l/ha.

Apricot, Cherry, Nectarines, Peach, Plum: 5 kg/ha. Commence applications at stone hardening and repeat at 10 to 14 day intervals up to one week before harvest. Water rate: 500-1000 l/ha.

Brassicas: 5 kg/ha. Commence application at 4 to 6 true leaves and repeat as necessary at 7 to 10 day intervals between applications. Water rate: 250-750 l/ha.

Carrots: 5 kg/ha. When sufficient leaf area to intercept spray with 10 to 14 day intervals between applications. Water rate: 200-500 l/ha.

Cereals: 5 kg/ha from mid-tillering to 2nd node detectable (Zadoks G.S. 25 to 32). Repeat at 10 to 14 day intervals between these growth stages as necessary. Water rate: 200 l/ha.

Citrus: 5 kg/ha. Commence applications after fruit set and repeat as necessary at 14 to 21 day intervals. Water rate: 1000 l/ha minimum.

Cucurbits (Field Grown): Apply 5 kg/ha at 10 day intervals from setting of first fruit / first fruit development. Water rate: 500 l/ha.

Grapevines: 5 kg/ha from fruit set and repeat at 10 to 14 day intervals as required until pre-harvest. Water rate: 500-1000 l/ha.

Kiwifruit: 5 kg/ha from 6 to 8 weeks post petal fall and repeated at 10 to 14 day intervals. Water rate: 500 to 1000 l/ha.

Onions: 5 kg/ha. When sufficient leaf area to intercept spray. Water rate: 200-500 l/ha.

Pears: 5 kg/ha at bud burst. Also, two to three applications at 10 to 14 day intervals, starting at petal fall. A further application of 5 kg/ha may be applied post-harvest, before leaf fall. Water rate: 500-1000 l/ha.

Potatoes: 5 kg/ha. Apply during bulking and repeat as necessary at 7 to 10 day intervals. Water rate: 200-500 l/ha.

Raspberry, Strawberry (Field Grown): 5 kg/ha. Commence applications after flowering and repeat as necessary at 7 to 10 day intervals. Water rate: 500-1000 l/ha.

Turf: 5 kg/ha. As soon as growth commences in the spring. In the case of moderate to severe deficiency, repeat applications at 14 day intervals. Water rate: 500-1000 l/ha.

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS
Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes (temperatures exceeding 40°C). Dispose of contents and container in accordance with all local, regional, national and international regulations

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.

