



Knowledge grows

# YaraVita™ CROPLIFT

A formulated multi-nutrient product for foliar application

Guaranteed Analysis

Total Nitrogen (N)	20%	Manganese (Mn)	0.26%
Phosphorus (P)	3.5%	Copper (Cu)	0.2%
Potassium (K)	11.6%	Zinc (Zn)	0.14%
Sulphur (S)	5%	Boron (B)	0.04%
Magnesium (Mg)	1.2%	Iron (Fe)	0.02%



## Why Foliar Apply?

Foliar sprays ensure precise application of the right nutrient mix at the right time, and can be specifically targeted to the leaf or fruit, to suit an immediate crop need.

Foliar application also provides nutrients for immediate uptake by the leaves or fruits. As a result, the grower is not reliant on the right soil, pH or growing media conditions and can quickly put the crop back on course

YaraVita CROPLIFT is a foliar fertiliser containing multiple nutrients and micronutrients for foliar application to a wide range of crops during times of stress or periods of rapid growth.

## Benefits:

- Formulated for safe application at critical growth stages to satisfy crop requirements.
- Widely tank mixable with other crop sprays. Visit [www.tankmix.com](http://www.tankmix.com) for details.
- Proven, reliable performance. Trialled and tested on a wide range of crops around the world
- High nutrient content means lower application rates reducing handling time and waste packaging.
- Easy to use soluble powder. Pours and disperses easily and quickly into the spray tank
- High quality, consistent products.
- Manufactured to ISO 9001 quality assurance standards.

Yara Fertilizers (NZ) Ltd

Tel: +64 6 8776600

Email: [nz.enquiries@yara.com](mailto:nz.enquiries@yara.com)

Website: [www.yara.co.nz](http://www.yara.co.nz)



## Application rates and timings

### Foliar Application

**All crops (field grown) except those specified below:** 2.5 to 5 kg/ha. Repeat at 10 to 14 day intervals as necessary. Avoid applications during flowering. Also, apply as necessary to perennial fruit crops post-harvest. Water rate: 200 l/ha (500 to 1000 l/ha for fruit crops). Alternatively: Spray to run off using a concentration of 2.5 grammes per litre. Allow 10 to 14 days between successive applications.

**Apple:** 5 kg/ha at 10 to 14 day intervals as required commencing 4 weeks after petal fall. Water rate: 500-1000 l/ha.

**Apricot, Cherry, Nectarines, Peach, Plum:** 5 kg/ha at 10 to 14 day intervals as required commencing at shuck fall. If necessary, also apply post-harvest before leaf fall. Water rate: 500 to 1000 l/ha.

**Brassicas:** 5 kg/ha. Commence application at 4 to 6 true leaves and repeat as necessary at 7 to 10 day intervals between applications. Water rate: 250-750 l/ha.

**Carrots:** 5 kg/ha. When sufficient leaf area to intercept spray with 10 to 14 day intervals between applications. Water rate: 200-500 l/ha.

**Cereals:** 5 kg/ha from mid-tillering to 2nd node detectable (Zadoks G.S. 25 to 32). Repeat at 10 to 14 day intervals between these growth stages as necessary. Water rate: 200 l/ha.

**Citrus:** 5 kg/ha applied during spring flush and also during summer/autumn flush. Water rate: 500 to 1000 l/ha.

**Cucurbits (Field Grown):** Apply 5 kg/ha at 10 day intervals from setting of first fruit / first fruit development. Water rate: 500 l/ha.

**Grapevines:** 2.5 to 5 kg/ha and repeat at 10 to 14 day intervals as required until pre-harvest. Avoid applications during flowering. Water rate: 500 to 1000 l/ha

**Kiwifruit:** 5 kg/ha and repeat at 10 to 14 day intervals as required until pre-harvest. Avoid applications during flowering. Water rate: 500 to 2000 l/ha. Water rate: 500 to 1000 l/ha.

**Onion:** 2.5 to 5 kg/ha. Repeat at 10 to 14 day intervals as necessary. Avoid applications during flowering. Water rate: 200 l/ha.

**Potatoes:** 2.5 to 5 kg/ha. Repeat at 10 to 14 day intervals as necessary. Avoid applications during flowering. Water rate: 200 l/ha.

**Raspberry, Strawberry (Field Grown):** 5 kg/ha. Commence applications after flowering and repeat as necessary at 7 to 10 day intervals. Water rate: 500-1000 l/ha.

**Turf:** 5 kg/ha. As soon as growth commences in the spring. In the case of moderate to severe deficiency, repeat applications at 14 day intervals. Water rate: 500-1000 l/ha.

## DIRECTIONS FOR USE

### ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

### TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit [www.tankmix.com](http://www.tankmix.com) for important information. Alternatively, contact Yara for specific advice

### PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

### STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes (temperatures exceeding 40°C). Dispose of contents and container in accordance with all local, regional, national and international regulations

### CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.

