

Growth stimulant with added nutrients for fruits and vegetables

SEAWEED FILTRATE GA142



Safety datasheet available on request









CROP RECOMMENDATIONS

Сгор	Rate/ Ha	No. of applications	Time of application	Directions for use : - shake well before use - pour Goëmar BM START la the tank Tank mixing and precautions : Goëmar BM START is compati with most usual fungicides at insecticides, but it is recommended to be mixed v herbicides. Storage : store the product normal conditions. Conditions of sales - lim warranty and limitations of liabi GOEMAR warrants that product within the unope container conforms to the chem description on this label. directions on this label must followed carefully and are ba on tests, believed to be reliable the occurrence of unusual weat or soil conditions, and the use this product, are all beyond control of the manufacturer, guarantee, expressed or impli is made as to the effects of s use or the results obtained, an no event shall the liability GOEMAR exceed the price of product used. The purchaser n assume all risks of storage, us application of this product. Go are sold subjectto the stanto Conditions of Sale of the manu turer, available on request. Us the product implies that purchaser accepts th Conditions of Sale.
Fruit trees - Apricots, peaches, cherries, nectarines, plums, olives, persimmons	2L	3	- Early flowering, petal fall, Young fruit stage	
Table and wine grapes	2 L	3	- Seperated flower buds - Full bloom - Early fruitset	
Cucurbits - Cucumber, melons, pumpkins, squash	2L	3	- First flower - 10-15 days after 1st application - 10-15 days after 2nd application	
Strawberries	2 L	3	- First flower - 10-15 days after 1st application - 10-15 days after 2nd application	
Solanaceae - Tomatoes, peppers, capsicums	2 L	3	- First flower - 10-15 days after 1st application - 10-15 days after 2nd application	
Leafy vegetables - Celery, lettuce, spinach, brassicas	2 L	3	- After planting out - 10-15 days after 1st application - 10-15 days after 2nd application	
Avocados	3 L	2	- Pre-flowering - Early fruitset	
Asparagus	2 L	2	- Beginning of branching - 10-15 days later	
Carrots	3 L	2	- 10 cm foliage - 10-15 days later	
Potatoes	2 L	3	- 1 week after tuber initiation - 7-14 days later - 7-14 days later	
Cereals	2 L	1	- Ear 1 cm to 2 nodes	
Peas - beans	2 L	2	- 15 cm stage - Beginning of flowering	

P102 : Keep out of the reach of children