



Knowledge grows

# YaraVita™ BRASSITREL PRO

A foliar multi-nutrient treatment for brassicas, legumes and other vegetable and arable crops

Guaranteed Analysis: total	w/v	
boron (B)	6%	60 g/L
manganese (Mn)	7.1%	71 g/L
molybdenum (Mo)	0.5%	5 g/L
magnesium (Mg)	7.1%	71 g/L
calcium (Ca)	8.9%	89 g/L
nitrogen (N)	6.9%	69 g/L

## Why Foliar Apply?

Foliar sprays ensure precise application of the right nutrient mix at the right time, and can be specifically targeted to the leaf or fruit, to suit an immediate crop need.

Foliar application also provides nutrients for immediate uptake by the leaves or fruits. As a result, the grower is not reliant on the right soil, pH or growing media conditions and can quickly put the crop back on course.

Magnesium, as well as the micronutrients boron, manganese and molybdenum are essential for crop establishment as well as winter hardiness of autumn sown crops. Stem extension is a period of high nutrient demand and often uptake of nutrients from the soil cannot keep pace with demand.

BRASSITREL PRO is formulated to supply all these vital nutrients in a concentrated, safe and balanced way at key moments in the growing season.



## Benefits:

- Formulated for safe application at critical growth stages to satisfy crop requirements
- Widely tank mixable with other crop sprays. Visit [www.tankmix.com](http://www.tankmix.com) for details.
- Proven, reliable performance. Trialled and tested on a wide range of crops around the world
- High quality, consistent product. Manufactured to ISO 9001 quality assurance standards
- Easy to use liquid formulation. Pours and disperses easily and quickly into the spray tank
- High nutrient content means lower application rates reducing handling time and waste packaging.



## Application Rates and Timings for YaraVita BRASSITREL PRO

**Beans, Legumes/Pulses, Peas, Soya Bean:** 3 to 4 l/ha as early in the growing season as possible, provided there is adequate leaf cover to intercept the spray (e.g. 4 to 6 leaf stage). Repeat as necessary at 10 to 14 day intervals up to one month before harvest.

N.B. Do not spray during flowering. Water rate: 200 l/ha.

**Brassicas, Broccoli, Brussel Sprouts, Cauliflower, Kale, Lettuce (Field Grown), Radish (Field Grown) Spinach, Swede, Turnip:** 3 to 4 l/ha as early in the growing season as possible, provided there is adequate leaf cover to intercept the spray (e.g. 4 to 6 leaf stage). Repeat as necessary at 10 to 14 day intervals up to one month before harvest. Water rate: 200 l/ha.

**Carrot:** Two applications of 3 l/ha applied when the crop is 15 cm tall and again 2 weeks later. Water rate: 200 l/ha.

**Cereals:** 3 l/ha applied at any time from the 2 leaf stage to 1st node detectable (Zadoks G.S. 12 to 31).

Water rate: 200 l/ha.

**Leek:** 3 l/ha applied two weeks after transplanting or in the case of direct sown crops, when the crop is 15 cm tall. For moderate to severe deficiency one or two repeat applications may be necessary at 10 to 14 day intervals. Water rate: 200 l/ha.

**Oilseed Rape:** 3 to 4 l/ha at the 4 to 9 leaf stage and again at the onset of the rapid stem extension phase up to the start of flowering. Repeat at 10 to 14 day intervals within these growth stages as necessary.

Water rate: 50 to 200 l/ha.

**Onion:** 2 applications of 3 l/ha applied as soon as there is adequate leaf cover and repeated 2 weeks later.

Water rate: 200 l/ha.

**Potatoes:** 3 l/ha applied one week after 100 % emergence with a second application 10 to 14 days later.

Water rate: 200 l/ha.

## Yara Crop Nutrition

Box 8746, Havelock North, New Zealand

Ph 06 877 6600, Fax 06 877 6610

Email: [nz.enquiries@yara.com](mailto:nz.enquiries@yara.com)

[www.yara.co.nz](http://www.yara.co.nz)

### LEGAL NOTICE

Yara International ASA and/or its group companies (collectively "Yara") make no express or implied warranty or representation concerning the accuracy or completeness of this document or the information contained in it. Information contained in this document is to the best of Yara's knowledge correct and accurate on the date of issuance. Any information provided is merely intended to serve as guidelines for the appropriate use, handling and storage of our products and may not be deemed as a guarantee or indication of quality, or serve as a basis for liability towards Yara in any way whatsoever. Any drawings, descriptive matter or illustrations contained in this document are provided for the sole purpose of giving an approximate idea of the products described in them. This document and any information contained in it shall remain the property of Yara. No rights, including, but not limited to, intellectual property rights, in respect of this document are granted to any recipient. Yara reserves the right to adjust and revise this document at any time. Please refer to our General Terms and Conditions for more information on legal matters.

